MOORE-VAN-ALLEN

# RECEIVED CENTRAL FAX CENTER SEP 1 0 2007

**2003/039** 

Docket No.: 020732-97.668 (7493) Appl. No. 10/792,038

# Section I. (The Claims)

- 1. (Currently Amended) A cleaning composition including a quaternary base, at least one alkali or alkaline earth base, and at least one additional component selected from the group consisting of a chelator, an oxirane species, and combinations thereof, wherein said chelator comprises a species selected from the group consisting of: 1-amino-1,2,4-triazole; 1-amino-1,2,3-triazole; 1-amino-5methyl-1,2,3-triazole; 3-amino-1,2,4-triazole; 3-mercapto-1,2,4-triazole; 3-isopropyl-1,2,4-triazole; naphthotriazole; 2-mercaptobenzoimidazole; 2-mercaptobenzothiazole; 5-aminotetrazole; 5-amino-1,3,4-thiadiazole-2-thiol; 2,4-diamino-6-methyl-1,3,5-triazine; thiazole; triazine; methyltetrazole; 1,3dimethyl-2-imidazolidinone: 1,5-pentamethylenetetrazole; 1-phenyl-5-mercaptotetrazole; diaminomethyltriazine; mercaptobenzothiazole; imidazoline thione; 4-methyl-4H-1,2,4-triazole-3thiol; 5-amino-1,3,4-thiadiazole-2-thiol; benzothiazole; tritolyl phosphate; indiazole; adenine; thioglycerol; salicylamide; iminodiacetic acid; benzoguanamine; melamine; thiocyranuric acid; anthranilic acid; 3-mercaptopropanol; and combinations thereof, and said cleaning composition is useful for removing photoresist and/or sacrificial anti-reflective coating (SARC) materials from a substrate having such material(s) thereon.
- 2. (Original) The cleaning composition of claim 1, which is devoid of hydroxylamine therein.
- 3. (Cancelled)
- 4. (Previously Presented) The cleaning composition of claim 1, comprising the following components:
  - 0.1 40.0 weight % organic quaternary base;
  - 0.01-5 weight % alkali or alkaline earth base;
  - 0-80 weight % solvent(s) and/or amine(s);
  - 0-5 weight % surfactant:
  - 0-10 weight % chelator/passivation agent; and
  - 0-98 weight % water.

wherein percentages of the components are percentages by weight, based on total weight of the composition, and wherein the total of the weight percentages of such components of the composition does not exceed 100 weight %.

5. (Previously Presented) The cleaning composition of claim 1, including at least one additional

Appl. No. 10/792,038

ingredient selected from the group consisting of stabilizers, dispersants, anti-oxidants, fillers, penetration agents, adjuvants, additives, and excipients.

- 6. (Previously Presented) The cleaning composition of claim 1, comprising the following components:
  - 2-15 weight % organic quaternary base;
  - ~0.01-2 weight % alkali or alkaline earth base;
  - 0-50 weight % solvent(s) and/or amine(s);
  - ~0.01-2 weight % surfactant;
  - 0-5 weight % chelator/passivation agent; and
  - 40 95 weight % water,

wherein percentages of the components are percentages by weight, based on total weight of the composition, and wherein the total of the weight percentages of such components of the composition does not exceed 100 weight %.

7. (Currently Amended) [[A]] The cleaning composition of claim 1, selected from the group consisting of Formulations A-C<sup>2</sup>, wherein all percentages are by weight, based on the total weight of the formulation:

## Formulation A

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

3.0% 4-methylmorpholine N-oxide

0.30% polyoxyethylene(150) dinonylphenyl ether

0.08% 2-mercaptobenzimidazole

91.0% water

#### Formulation B

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

3.0% 4-methylmorpholine N-oxide

0.30% polyoxyethylene(150) dinonylphenyl ether

0.20% 5-amino-1,3,4-thiadiazole-2-thiol

90.86% water

## Formulation C

3.60% benzyltrimethylammonium hydroxide

0.27% potassium hydroxide

3.5% 4-methylmorpholine N-oxide

15.0% 4-(3-aminopropyl)morpholine 0.30% polyoxyethylene(150) dinonylphenyl ether 0.08% 2-mercaptobenzimidazole 77.25% water

#### Formulation D

5.36% benzyltrimethylammonium hydroxide 0.28% potassium hydroxide 20.0% dimethyl sulfoxide 0.08% 2-mercaptobenzimidazole 74.28% water

#### Formulation E

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

10.0% tetramethylene sulfone

0.30% oxirane, methyl-, polymer with oxirane, ether with 2,2'-(oxidoimino)bis(ethanol) (2:1), N(-3(C9-11-isoalkyloxy)propyl)derivatives, C<sub>10</sub>-rich

0.08% 2-mercaptobenzimidazole

83.98% water

#### Formulation F

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

10.0% di(ethyleneglycol)butyl ether

10.0% 2-(2-dimethylamino)ethoxy)ethanol

0.30% oxirane, methyl-, polymer with oxirane, ether with 2,2'-(oxidoimino)bis(ethanol) (2:1), N(-3(C9-11-isoalkyloxy)propyl)derivatives, C<sub>10</sub>-rich

74.06% water

# Formulation G

5.36% benzyltrimeth	ylammonium	hydroxide
---------------------	------------	-----------

0.28% potassium hydroxide

10.0% tetramethylene sulfone

10.0% di(ethyleneglycol)butyl ether

0.10% oxirane, methyl-, polymer with oxirane, mono(octylphenyl)ether

0.08% 2-mercaptobenzimidazole

74.18% water,

## Formulation H

benzyltrimethylammonium hydroxide, 40% aqueous solution	9.0 %
potassium hydroxide, 45% aqueous solution	0.6 %
N-methylmorpholine oxide, 50% aqueous solution dinonylphenol ethoxylate, 7% aqueous solution	<b>~7.0 %</b>
dinonylphenol ethoxylate, 7% aqueous solution	4.3 %
2-mercaptobenzimidazole	0.1 %
aminopropylmorpholine	20.0 %

MOORE-VAN-ALLEN

	Docket No.: 020732-97.668 (7493) Appl. No. 10/792,038
water	59.02 %
Formulation I	·
benzyltrimethylammonium hydroxide, 40% aqueous solution potassium hydroxide, 45% aqueous solution N-methylmorpholine oxide, 50% aqueous solution dinonylphenol ethoxylate, 7% aqueous solution 2-mercaptobenzimidazole aminopropylmorpholine water	9.0 % 0.6 % 7.0 % 4.3 % 0.1 % 15.0 %
water	64.02 %
Formulation J	·
benzyltrimethylammonium hydroxide, 40% aqueous solution potassium hydroxide, 45% aqueous solution N-methylmorpholine oxide, 50% aqueous solution dinonylphenol ethoxylate, 7% aqueous solution 2-mercaptobenzimidazole aminopropylmorpholine water	9.0 % 0.6 % 7.0 % 4.3 % 0.1 % 10.0 % 69.02 %
Formulation K	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	13.4 % 7.0 % 0.6 % 0.08 % 0.3 % 78.62 %
Formulation L	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	13.4 % 7.0 % 1.2 % 0.08 % 0.3 % 78.02 %
Formulation M	
tetramethylammonium hydroxide, 25% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	5.85 % 7.0 % 1.2 % 0.08 % 0.3 % 85.57 %

# Formulation N

	·
	Docket No.: 020732-97.668 (7493)
	Appl. No. 10/792,038
totromathylammanium hudrovide 250/	2.02.84
tetramethylammonium hydroxide, 25% aqueous solution	2.93 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
KOH, 45% aqueous solution	1.2 %
2-mercaptobenzimidizole	0.08 %
dinonylphenol polyoxyethylene	0.3 %
water	88.49 %
Formulation O	
benzyltrimethylammonium hydroxide, 40% aqueous solution	. 7.2 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
KOH, 45% aqueous solution	0.6 %
2-mercaptobenzimidizole	
dinonylphenol polyoxyethylene	0.08 %
water	0.3 %
water	84.82 %
Formulation P	
benzyltrimethylammonium hydroxide, 40% aqueous solution	3.6 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
KOH, 45% aqueous solution	1.2 %
2-mercaptobenzimidizole	0.08 %
dinonylphenol polyoxyethylene	·
water	0.3 %
Water	87.82 %
Formulation Q	
benzyltrimethylammonium hydroxide, 40% aqueous solution	3.6 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
KOH, 45% aqueous solution	0.6 %
2-mercaptobenzimidizole	0.08 %
dinonylphenol polyoxyethylene	0.3 %
water	88.42 %
	00.72 /0
Formulation R	
benzyltrimethylammonium hydroxide, 40% aqueous solution	7.2 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
KOH, 45% aqueous solution	0.3 %
2-mercaptobenzimidizole	0.08 %
dinonylphenol polyoxyethylene	0.3 %
water	85.12 %,
Formulation S	
benzyltrimethylammonium hydroxide, 40% aqueous solution	22.26 %
Potassium hydroxide, 45% aqueous solution	0.6 %
2-mercaptobenzimidazole	
methyldiethanolamine	2.33 %
phosphoric acid (86 %)	1.69 %
2 omino 6 managente 1.0 4 tuismala	1.07 /0

1.0 %

3-amino-5-mercapto-1,2,4-triazole

Formulation Y

Docket No.: 020732-97.668 (7493) Appl. No. 10/792,038

water	72.04 %
Formulation T	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 4-methyl-2-phenyl-imidazole water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation U	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 2-mercaptothiazoline water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation V	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 8-hydroxyquinoline water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation W	•
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86%) 1-phenyl-2-tetrazoline-5-thione water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation X	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) gallic acid water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %

	Docket No.: 020732-97.668 (7493) Appl. No. 10/792,038
benzyltrimethylammonium hydroxide, 40% aqueous solution	22.26 %
Potassium hydroxide, 45% aqueous solution	0.6 %
2-mercaptobenzimidazole	0.08 %
methyldiethanolamine	2.33 %
phosphoric acid (86 %)	1.69 %
salicylic acid	1.0 %
water	72.04 %
	72.04 78
Formulation Z	
benzyltrimethylammonium hydroxide, 40% aqueous solution	22.26 %
Potassium hydroxide, 45% aqueous solution	0.6 %
2-mercaptobenzimidazole	0.08 %
methyldiethanolamine	2.33 %
phosphoric acid (86 %)	1.69 %
ascorbic acid	1.0 %
water	72.04 %
_	
Formulation A <sup>2</sup>	
benzyltrimethylammonium hydroxide, 40% aqueous solution	7.2 %
Potassium hydroxide, 45% aqueous solution	0.6 %
2-mercaptobenzimidazole	0.08 %
aminopropyl morpholine	10 %
4-methyl-2-phenyl-imidazole	1.0 %
water	81.12 %
	01.12 70
Formulation B <sup>2</sup>	
benzyltrimethylammonium hydroxide, 40% aqueous solution	7.2 %
Potassium hydroxide, 45% aqueous solution	0.6 %
2-mercaptobenzimidazole	0.08 %
aminopropyl morpholine	10 %
4-methyl-2-phenyl-imidazole	0.5 %
water	81.62 %
Formulation C <sup>2</sup>	
benzyltrimethylammonium hydroxide, 40% aqueous solution	7.2 %
Potassium hydroxide, 45% aqueous solution	0.6 %
2-mercaptobenzimidazole	0.08 %
aminopropyl morpholine	10 %
4-methyl-2-phenyl-imidazole	1.0 %
water	81.02 %
dinonylphenol polyoxyethylene	0.1 %
	V+1 /U

wherein said cleaning composition is useful for removing photoresist and/or sacrificial anti-reflective coating (SARC) materials from a substrate having such material(s) thereon.

# 8. (Cancelled)

Appl. No. 10/792,038

- 9. (Cancelled)
- 10. (Previously Presented) The cleaning composition of claim 1, wherein the alkali base comprises potassium hydroxide.
- 11.-13. (Cancelled)
- 14. (Previously Presented) The cleaning composition of claim 1, wherein the chelator comprises 2-mercaptobenzimidazole.
- 15. (Previously Presented) The cleaning composition of claim 1, wherein the chelator is present in an amount greater than about 0.08 wt.%, based on the total weight of the composition.
- 16. (Cancelled)
- 17. (Previously Presented) The composition of claim 1, further comprising a surfactant.
- 18. (Original) The composition of claim 17, wherein the surfactant comprises a surfactant species selected from the group consisting of: fluoroalkyl surfactants; polyethylene glycols; polypropylene glycols; polyethylene glycol ethers; polypropylene glycol ethers; carboxylic acid salts; dodecylbenzenesulfonic acid and salts thereof; polyacrylate polymers; dinonylphenyl polyoxyethylene; silicone polymers; modified silicone polymers; acetylenic diols; modified acetylenic diols, alkylammonium salts; modified alkylammonium salts; and combinations of two or more of the foregoing.
- 19. (Previously Presented) The composition of claim 1, further comprising a co-solvent.
- 20. (Original) The composition of claim 19, wherein the co-solvent comprises a co-solvent species selected from the group consisting of: amines; glycols; glycol ethers; polyglycol ethers; and combinations of two or more of the foregoing.
- 21. (Original) The composition of claim 19, wherein the co-solvent comprises a co-solvent species selected from the group consisting of: dimethyldiglycolamine; 1,8-diazabicyclo[5.4.0]undecene;

aminopropylmorpholine; triethanolamine; methylethanolamine; diethylene glycol; propylene glycol; neopentyl glycol; hydroxyethylmorpholine; aminopropylmorpholine; di(ethylene glycol)monoethyl ether; di(propylene glycol)propyl ether; ethylene glycol phenyl ether; di(propylene glycol) butyl ether; butyl carbitol; polyglycol ethers; and combinations of two or more of the foregoing.

- 22. (Cancelled)
- 23. (Cancelled)
- 24. (Withdrawn) A method of removing photoresist and/or SARC material from a substrate having said material thereon, said method comprising contacting the substrate with a cleaning composition for sufficient time to at least partially remove said material from the substrate, wherein the cleaning composition includes a quaternary base, at least one alkali or alkaline earth base, and at least one additional component selected from the group consisting of a chelator, an oxirane species, and combinations thereof, wherein said chelator comprises a species selected from the group consisting of: 1-amino-1,2,4-triazole; 1-amino-1,2,3-triazole; 1-amino-5-methyl-1,2,3-triazole; 3amino-1,2,4-triazole; 3-mercapto-1,2,4-triazole; 3-isopropyl-1,2,4-triazole; naphthotriazole; 2mercaptobenzoimidazole; 2-mercaptobenzothiazole; 5-aminotetrazole; 5-amino-1,3,4-thiadiazole-2thiol; 2,4-diamino-6-methyl-1,3,5-triazine; thiazole; triazine; methyltetrazole; 1,3-dimethyl-2imidazolidinone; 1,5-pentamethylenetetrazole; 1-phenyl-5-mercaptotetrazole; diaminomethyltriazine; mercaptobenzothiazole; imidazoline thione; 4-methyl-411-1,2,4-triazole-3-thiol; 5-amino-1,3,4thiadiazole-2-thiol; benzothiazole; tritolyl phosphate; indiazole; adenine; thioglycerol; salicylamide; iminodiacetic acid; benzoguanamine; melamine; thiocyranuric acid; anthranilic acid; 3mercaptopropanol; and combinations thereof.
- 25. (Withdrawn) The method of claim 24, wherein the substrate comprises a semiconductor device structure.
- 26. (Withdrawn) The method of claim 24, wherein the material comprises photoresist.

WATER TO THE TANK OF THE PARTY OF THE PARTY

- 27. (Withdrawn) The method of claim 24, wherein the material comprises SARC material.
- 28. (Withdrawn) The method of claim 27, wherein the SARC material has been applied to a semiconductor device structure to minimize reflectivity variations during photolithographic patterning

Appl. No. 10/792,038

on the semiconductor device structure.

- 29. (Withdrawn) The method of claim 24, wherein said contacting is carried out for a time of from about 10 to about 45 minutes.
- 30. (Withdrawn) The method of claim 24, wherein said contacting is carried out at temperature in a range of from about 50°C to about 80°C.
- 31. (Withdrawn) The method of claim 24, wherein the composition is devoid of hydroxylamine therein.
- 32. (Cancelled)
- 33. (Withdrawn) The method of claim 24, wherein the composition comprises the following components:
  - 0.1 40.0 weight % organic quaternary base;
  - 0.01-5 weight % alkali or alkaline earth base;
  - 0-80 weight % solvent(s) and/or amine(s);
  - 0-5 weight % surfactant:
  - 0-10 weight % chelator/passivation agent; and
  - 0-98 weight % water,

wherein percentages of the components are percentages by weight, based on total weight of the composition, and wherein the total of the weight percentages of such components of the composition does not exceed 100 weight %.

- 34. (Withdrawn) The method of claim 24, wherein the composition includes at least one additional ingredient selected from the group consisting of stabilizers, dispersants, anti-oxidants, fillers, penetration agents, adjuvants, additives, fillers, and excipients.
- 35. (Withdrawn) The method of claim 24, wherein the composition comprises the following components:
- 2-15 weight worganic quaternary base;
  - ~0.01-2 weight % alkali or alkaline earth base;
  - 0-50 weight % solvent(s) and/or amine(s);

Appl. No. 10/792,038

- ~0.01-2 weight % surfactant;
- 0-5 weight % chelator/passivation agent; and
- 40-95 weight % water,

wherein percentages of the components are percentages by weight, based on total weight of the composition, and wherein the total of the weight percentages of such components of the composition does not exceed 100 weight %.

36. (Withdrawn) [[A]] The method of claim 24 of removing photoresist and/or SARC material from a substrate having said material thereon, said method comprising contacting the substrate with a cleaning composition for sufficient time to at least partially remove said material from the substrate, wherein the cleaning composition is selected from the group consisting of Formulations A-C<sup>2</sup>, wherein all percentages are by weight, based on the total weight of the formulation:

#### Formulation A

- 5.36% benzyltrimethylammonium hydroxide
- 0.28% potassium hydroxide
- 3.0% 4-methylmorpholine N-oxide
- 0.30% polyoxyethylene(150) dinonylphenyl ether
- 0.08% 2-mercaptobenzimidazole
- 91.0% water

# Formulation B

- 5.36% benzyltrimethylammonium hydroxide
- 0.28% potassium hydroxide
- 3.0% 4-methylmorpholine N-oxide
- 0.30% polyoxyethylene(150) dinonylphenyl ether
- 0.20% 5-amino-1,3,4-thiadiazole-2-thiol
- 90.86% water

# Formulation C

- 3.60% benzyltrimethylammonium hydroxide
- 0.27% potassium hydroxide
- 3.5% 4-methylmorpholine N-oxide
- 15.0% 4-(3-aminopropyl)morpholine
- 0.30% polyoxyethylene(150) dinonylphenyl ether
- 0.08% 2-mercaptobenzimidazole
- 77.25% water

#### Formulation D

- 5.36% benzyltrimethylammonium hydroxide
- 0.28% potassium hydroxide

20.0% dimethyl sulfoxide 0.08% 2-mercaptobenzimidazole 74.28% water

# Formulation E

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

10.0% tetramethylene sulfone

0.30% oxirane, methyl-, polymer with oxirane, ether with 2,2'-(oxidoimino)bis(ethanol) (2:1), N(-3(C9-11-isoalkyloxy)propyl)derivatives, C<sub>10</sub>-rich

0.08% 2-mercaptobenzimidazole

83.98% water

## Formulation F

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

10.0% di(ethyleneglycol)butyl ether

10.0% 2-(2-dimethylamino)ethoxy)ethanol

0.30% oxirane, methyl-, polymer with oxirane, ether with 2,2'-(oxidoimino)bis(ethanol) (2:1), N(-3(C9-11-isoalkyloxy)propyl)derivatives, C<sub>10</sub>-rich

74.06% water

## Formulation G

5.36% benzyltrimethylammonium hydroxide

0.28% potassium hydroxide

10.0% tetramethylene sulfone

10.0% di(ethyleneglycol)butyl ether

0.10% oxirane, methyl-, polymer with oxirane, mono(octylphenyl)ether

0.08% 2-mercaptobenzimidazole

74.18% water,

## Formulation H

benzyltrimethylammonium hydroxide, 40% aqueous solution	9.0 %
potassium hydroxide, 45% aqueous solution	0.6 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
dinonylphenol ethoxylate, 7% aqueous solution	4.3 %
2-mercaptobenzimidazole	0.1 %
aminopropylmorpholine	20.0 %
water	59.02 %

# Formulation I

benzyltrimethylammonium hydroxide, 40% aqueous solution	9.0 %
potassium hydroxide, 45% aqueous solution	0.6 %
N-methylmorpholine oxide, 50% aqueous solution	7.0 %
dinonylphenol ethoxylate, 7% aqueous solution	4.3 %
2-mercaptobenzimidazole	0.1 %

	Docket No.: 020732-97.668 (7493) Appl. No. 10/792,038
aminopropylmorpholine water	15.0 % 64.02 %
Formulation J	
benzyltrimethylammonium hydroxide, 40% aqueous solution potassium hydroxide, 45% aqueous solution N-methylmorpholine oxide, 50% aqueous solution dinonylphenol ethoxylate, 7% aqueous solution 2-mercaptobenzimidazole aminopropylmorpholine water	9.0 % 0.6 % 7.0 % 4.3 % 0.1 % 10.0 % 69.02 %
Formulation K	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	13.4 % 7.0 % 0.6 % 0.08 % 0.3 % 78.62 %
Formulation L	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	13.4 % 7.0 % 1.2 % 0.08 % 0.3 % 78.02 %
Formulation M	
tetramethylammonium hydroxide, 25% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	5.85 % 7.0 % 1.2 % 0.08 % 0.3 % 85.57 %
Formulation N	
tetramethylammonium hydroxide, 25% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	2.93 % 7.0 % 1.2 % 0.08 % 0.3 % 88.49 %

# Formulation O

	Docket No.: 020732-97.668 (7493) Appl. No. 10/792,038
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	7.2 % 7.0 % 0.6 % 0.08 % 0.3 % 84.82 %
Formulation P	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	3.6 % 7.0 % 1.2 % 0.08 % 0.3 % 87.82 %
Formulation Q	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	3.6 % 7.0 % 0.6 % 0.08 % 0.3 %
Formulation R	
benzyltrimethylammonium hydroxide, 40% aqueous solution N-methylmorpholine oxide, 50% aqueous solution KOH, 45% aqueous solution 2-mercaptobenzimidizole dinonylphenol polyoxyethylene water	7.2 % 7.0 % 0.3 % 0.08 % 0.3 % 85.12 %,
Formulation S	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 3-amino-5-mercapto-1,2,4-triazole water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation T	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 4-methyl-2-phenyl-imidazole	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 %

Docket No.: 020732-9	7.668 (7493)
Appl. No	. 10/792,038

water	72.04 %
Formulation U	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 2-mercaptothiazoline water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation V	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) 8-hydroxyquinoline water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation W	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86%) 1-phenyl-2-tetrazoline-5-thione water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation X	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) gallic acid water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation Y	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) salicylic acid water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %

# Formulation Z

benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole methyldiethanolamine phosphoric acid (86 %) ascorbic acid water	22.26 % 0.6 % 0.08 % 2.33 % 1.69 % 1.0 % 72.04 %
Formulation A <sup>2</sup>	, 2.0 . , 0
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole aminopropyl morpholine 4-methyl-2-phenyl-imidazole water	7.2 % 0.6 % 0.08 % 10 % 1.0 % 81.12 %
Formulation B <sup>2</sup>	•
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole aminopropyl morpholine 4-methyl-2-phenyl-imidazole water	7.2 % 0.6 % 0.08 % 10 % 0.5 % 81.62 %
Formulation C <sup>2</sup>	
benzyltrimethylammonium hydroxide, 40% aqueous solution Potassium hydroxide, 45% aqueous solution 2-mercaptobenzimidazole aminopropyl morpholine 4-methyl-2-phenyl-imidazole water dinonylphenol polyoxyethylene	7.2 % 0.6 % 0.08 % 10 % 1.0 % 81.02 % 0.1 %.

- 37. (Cancelled)
- 38. (Cancelled)
- 39. (Withdrawn) The method of claim 24, wherein the alkali base comprises potassium hydroxide.
- 40.-42. (Cancelled)
- 43. (Withdrawn) The method of claim 24, wherein the chelator comprises 2-mercaptobenzimidazole.

44. (Withdrawn) The method of claim 43, wherein the chelator is present in an amount greater than about 0.08 wt.%, based on the total weight of the composition.

- 45. (Withdrawn) The method of claim 24, wherein the quaternary base comprises benzyltrimethylammonium hydroxide.
- 46. (Withdrawn) The method of claim 24, wherein the cleaning composition further comprises a surfactant.
- 47. (Withdrawn) The method of claim 46, wherein the surfactant comprises a surfactant species selected from the group consisting of: fluoroalkyl surfactants; polyethylene glycols; polypropylene glycols; polyethylene glycol ethers; polypropylene glycol ethers; carboxylic acid salts; dodecylbenzenesulfonic acid and salts thereof; polyacrylate polymers; dinonylphenyl polyoxyethylene; silicone polymers; modified silicone polymers; acetylenic diols; modified acetylenic diols, alkylammonium salts; modified alkylammonium salts; and combinations of two or more of the foregoing.
- 48. (Withdrawn) The method of claim 24, wherein the cleaning composition further comprises a co-solvent.
- 49. (Withdrawn) The method of claim 48, wherein the co-solvent comprises a co-solvent species selected from the group consisting of: amines; glycols; glycol ethers; polyglycol ethers; and combinations of two or more of the foregoing.
- 50. (Withdrawn) The method of claim 48, wherein the co-solvent comprises a co-solvent species selected from the group consisting of: dimethyldiglycolamine; 1,8-diazabicyclo[5.4.0]undecene; aminopropylmorpholine; triethanolamine; methylethanolamine; diethylene glycol; propylene glycol; neopentyl glycol; hydroxyethylmorpholine; aminopropylmorpholine; di(ethylene glycol)monoethyl ether; di(propylene glycol)propyl ether; ethylene glycol phenyl ether; di(propylene glycol) butyl ether; butyl carbitol; polyglycol ethers; and combinations of two or more of the foregoing.
- 51. (Cancelled)
- 52. (Cancelled)

- 53. (Previously Presented) The cleaning composition of claim 1, wherein the quaternary base comprises an organic quaternary ammonium base.
- 54. (Previously Presented) The cleaning composition of claim 1, wherein the quaternary base comprises benzyltrimethylammonium hydroxide.
- 55. (Previously Presented) The cleaning composition of claim 1, wherein the ACC comprises benzyltrimethylammonium hydroxide and potassium hydroxide.
- 56. (Previously Presented) The cleaning composition of claim 19, wherein the co-solvent comprises a glycol ether.
- 57. (Previously Presented) The cleaning composition of claim 1 comprising benzyltrimethylammonium hydroxide; potassium hydroxide; tetramethylene sulfone; di(ethyleneglycol)butyl ether; oxirane, methyl-, polymer with oxirane, mono(octylphenyl)ether; 2-mercaptobenzimidazole; and water.
- 58. (Withdrawn) A method of making a semiconductor device comprising contacting the substrate with the cleaning composition of claim 1 for sufficient time to at least partially remove said material from the substrate.
- 59. (Previously Presented) The cleaning composition of claim 1, further comprising oxidant.